CPEB4

CONTACT INFORMATION: Monoclonal Antibodies Unit. Centro Nacional de Investigaciones Oncológicas

STATUS: Validated

TYPE: mouse anti-mouse

CLONE NAME: ERE93C

PROTEIN: CPEB4 (Cytoplasmic polyadenylation element-binding protein 4)

PROTEIN WEB: http://www.uniprot.org/uniprot/Q7TN98

ANTIGEN USED: HIS-GST-mCPEB4 (1-375aa) recombinant protein

FUSION PARTNER: NS1/Ag4-1 (NS1) cells

ISOTYPE: IgG1

SPECIES REACTIVITY: human and mouse

PREPARATION AND STORAGE: Aliquot and store at 4C. Do not freeze

PUBLICATION DESCRIBING ANTIBODY CHARACTERIZATION/VALIDATION

Lineage-specific roles of the cytoplasmic polyadenylation factor CPEB4 in the regulation of melanoma drivers. Pérez-Guijarro E, Karras P, Cifdaloz M, Martínez-Herranz R, Cañón E, Graña O, Horcajada-Reales C, Alonso-Curbelo D, Calvo TG, Gómez-López G, Bellora N, Riveiro-Falkenbach E, Ortiz-Romero PL, Rodríguez-Peralto JL, Maestre L, Roncador G, de Agustín Asensio JC, Goding CR, Eyras E, Megías D, Méndez R, Soengas MS.


Nat Commun. 2016 Nov 18; 7:13418.

APPLICATIONS

IHC Techniques	Clone	Dilution	Antibody concentration	Antigen retrieval method	Visualization kit	Positive control	Negative control	Protein localization	Positivity in other species	
Frozen tissue and cytospins										
Recommended	Neat	Supernat								
		ant								
Paraffin tissue										


Monoclonal Antibodies Catalogue

Recommended	1:1500	Purified							
Immunofluorescence									
Recommended	Neat	Supernat							
		ant							



ERE93C mAb in transfected cells

To confirm that ERE93C recognizes mCPEB4 immunohistochemistry on frozen cytospins preparations of GFP or Flag-tagged mCPEB1, mCPEB3 and mCPEB4 expressed in Hek293T was performed. Anti-Flag and anti-GFP antibodies were used as positive controls.




ERE93C staining in mouse paraffin sections.

ERE93C antibody can be used to detect mCPEB4 protein in mouse paraffin embedded tissue samples.

Tissue sample: mouse brain and mouse embryonic tissues.

Dilution: 1:1500

Antigen retrieval: CC1 D5+OR
Antibody incubation: 30 minutes
Detection kit used: Ventana



ERE93C antibody can be used to detect mCPEB4 protein by Immunofluorescence.

Tissue sample: Mefs.