

## P16


**CONTACT INFORMATION:** Monoclonal Antibodies Unit. Centro Nacional de Investigaciones Oncológicas  
**STATUS:** Validated  
**TYPE:** rat anti mouse  
**CLONE NAME:** PABLO33B  
**PROTEIN:** cyclin-dependent kinase inhibitor 2A  
**ANTIGEN USED:** HIS-GST-mp16 recombinant protein (full length protein)  
**FUSION PARTNER:** myeloma p3-NS1/Ag4-1 (NS1) cells  
**ISOTYPE:** IgG2a  
**SPECIES REACTIVITY:** mouse  
**PREPARATION AND STORAGE:** Aliquot and store at 4C. Do not freeze

### REFERENCES

Therapeutic senescence via GPCR activation in synovial fibroblasts facilitates resolution of arthritis. Trinidad Montero-Melendez, Ai Nagano, Claude Chelala, Andrew Filer, Christopher D. Buckley and Mauro Perretti. Nature Communications volume 11, Article number: 745 (2020)


### APPLICATIONS

IHC Techniques	Clone	Dilution	Antibody concentration	Antigen retrieval method	Visualization kit	Positive control	Negative control	Protein localization	Positivity in other species
<b>Frozen tissue and cytopins</b>									
Recommended	PABL O33B	Neat	supernatant						
<b>Paraffin tissue</b>									
Recommended	PABL O33B	Neat	supernatant		Ventana	Fibrosarcoma tissue		nuclear	
<b>Immunofluorescence</b>									




**PABLO33B is able to detect mouse p16 (CDKN2A) protein in immunocytochemistry**

To confirm that PABLO33B mAb recognizes mouse p16 (CDKN2A) protein, immunocytochemistry on frozen cytopins preparations of HEK-mp16-V5 transfected cells was performed. Cytopsin preparation of V5-tagged mouse p15 protein was used as a negative control. Anti-V5 was used as a positive control.



**PABLO33B mAb can be used to detect mouse p16 (CDKN2A) protein in paraffin tissues**

WB Techniques	Clone	Dilution	Antibody concentration	Positive control	Negative control	Expected MW	Observed Mw	Positivity in other species
<b>Western Blotting</b>								
Recommended	PABLO33B	undiluted	supernatant			16kDa	16kDa	
<b>Immunoprecipitation</b>								




## PABLO33B

**PABLO33B mAb is able to detect mouse p16 (CDKN2A) protein by WB.**

LANES

Lane 1 HEKmp16 (20ug) (+)

Lane 2 HEKmp15 (20ug) (-)



## PABLO33B

**PABLO33B mAb is able to detect mouse p16 (CDKN2A) protein by WB.**

LANES

Lane 1 NS1 mouse cell line (100ug)

Lane 2 PAP1 mouse papilloma tissue (100ug)

Lane 3 PAP3 mouse papilloma tissue (100ug)